

## SIS and SIR Epidemic Models Under Virtual Dispersal

Derdei Bichara<sup>1</sup> · Yun Kang<sup>2</sup> · Carlos Castillo-Chavez<sup>1</sup> ·  
Richard Horan<sup>3</sup> · Charles Perrings<sup>4</sup>

Received: 18 February 2015 / Accepted: 2 October 2015  
© Society for Mathematical Biology 2015

**Abstract** We develop a multi-group epidemic framework via virtual dispersal where the risk of infection is a function of the residence time and local environmental risk. This novel approach eliminates the need to define and measure contact rates that are used in the traditional multi-group epidemic models with heterogeneous mixing. We apply this approach to a general  $n$ -patch SIS model whose basic reproduction number  $\mathcal{R}_0$  is computed as a function of a patch residence-time matrix  $\mathbb{P}$ . Our analysis implies that the resulting  $n$ -patch SIS model has robust dynamics when patches are strongly connected: There is a unique globally stable endemic equilibrium when  $\mathcal{R}_0 > 1$ , while the disease-free equilibrium is globally stable when  $\mathcal{R}_0 \leq 1$ . Our further analy-

---

✉ Yun Kang  
yun.kang@asu.edu

Derdei Bichara  
derdei.bichara@asu.edu

Carlos Castillo-Chavez  
ccchavez@asu.edu

Richard Horan  
horan@msu.edu

Charles Perrings  
Charles.Perrings@asu.edu

- <sup>1</sup> SAL Mathematical, Computational and Modeling Science Center, Arizona State University, Tempe, AZ 85287, USA
- <sup>2</sup> Sciences and Mathematics Faculty, College of Letters and Sciences, Arizona State University, Mesa, AZ 85212, USA
- <sup>3</sup> Department of Agricultural, Food and Resource Economics, Michigan State University, East Lansing, MI 48824, USA
- <sup>4</sup> School of Life Sciences, Arizona State University, Tempe, AZ 85287, USA

sis indicates that the dispersal behavior described by the residence-time matrix  $\mathbb{P}$  has profound effects on the disease dynamics at the single patch level with consequences that proper dispersal behavior along with the local environmental risk can either promote or eliminate the endemic in particular patches. Our work highlights the impact of residence-time matrix if the patches are not strongly connected. Our framework can be generalized in other endemic and disease outbreak models. As an illustration, we apply our framework to a two-patch SIR single-outbreak epidemic model where the process of disease invasion is connected to the final epidemic size relationship. We also explore the impact of disease-prevalence-driven decision using a phenomenological modeling approach in order to contrast the role of constant versus state-dependent  $\mathbb{P}$  on disease dynamics.

**Keywords** Epidemiology · SIS–SIR models · Dispersal · Residence times · Global stability · Adaptive behavior · Final size relationship

**Mathematics Subject Classification** Primary 34D23 · 92D25 · 60K35

## 1 Introduction

Sir Ronald Ross must be considered the founder of mathematical epidemiology (Ross 1911) despite the fact that Daniel Bernoulli (1700–1782) was most likely the first researcher to introduce the use of mathematical models in the study of epidemic outbreaks (Bernoulli 1766; Dietz and Heesterbeek 2002) nearly 150 years earlier. Ross appendix to his 1911 paper (Ross 1911) not only introduces a nonlinear system of differential equations aimed at capturing the overall dynamics of the malaria contagion, a disease driven by the interactions of hosts, vectors and the life history of *Plasmodium falciparum*, but also includes a tribute to mathematics through his observation that this framework may also be used to model the dynamics of sexually transmitted diseases (Ross 1911). Ross observation has motivated the use of mathematics in the study of the impact of human social interaction on disease dynamics (Blythe and Castillo-Chavez 1989; Castillo-Chavez and Busenberg 1991; Castillo-Chavez and Huang 1999; Castillo-Chavez et al. 1996, 1999; Haderler and Castillo-Chavez 1995; Hethcote and Yorke 1984; Hsu Schmitz 2000a, b, 2007; Yorke et al. 1978). In particular, Ross work introduced the type of framework needed to capture and modify the dynamics of epidemic outbreaks: new landscapes where public policies could be tried and tested without harming anybody, complementing and expanding the role that statistics plays in epidemiology. Suddenly scientists and public health experts had a “laboratory” for assessing the impact of transmission mechanisms, evaluating, a priori, efforts aimed at mitigating or eliminating the deleterious impact of disease dynamics.

The study of the dynamics of communicable disease in metapopulation, multi-group or age-structure models has also benefitted from the work of Ross. Contact matrices have been used in the study of disease dynamics to accommodate or capture the dynamics of heterogeneous mixing populations (Anderson and May 1982; Castillo-Chavez et al. 1989; Dietz and Schenzle 1985; Hethcote 2000). The spread

of communicable diseases like measles, chicken pox or rubella is intimately connected to the concept of contact, “effective” contact or “effective” per capita contact rate (Castillo-Chavez et al. 1994; Hethcote 2000), a clear measurable concept in, for example, the context of sexually transmitted diseases (STDs) or vector-borne diseases. The values used to define a contact matrix emerge from the a priori belief that contacts can be clearly defined and measured in any context. Their use in the context of communicable diseases is based often on relative rankings, the result of observational subjective measures of contact or activity levels. For example, since children are believed to have the most contacts per unit of time, their observed activity levels are routinely used to set a relative contact or activity scale. Traditionally, since school children are assumed to be the most active, they are used to set the scale with the rest of the age-specific contact matrix usually completed under the assumption of proportionate (weighted random) mixing (albeit other forms of mixing are possible Anderson and May 1991; Blythe and Castillo-Chavez 1989; Castillo-Chavez and Busenberg 1991; Castillo-Chavez et al. 1989; Dietz and Schenzle 1985; Hethcote 2000 and references therein). In short, mixing or contact matrices are used to collect rescaled estimated levels of activity among interacting subgroups or age-classes, a phenomenological estimation process based on observational studies and surveys (Mossong et al. 2008). Our belief that contact rates cannot, in general, be measured in satisfactory ways for diseases like influenza, measles or tuberculosis arises from the difficulty of assessing the average number of contacts per unit of time of children in a school bus or the average number of contacts per unit of time that children and adults have with each other in a classroom or at the library, per unit of time. The issue is further confounded by our inability to assess what an effective contact is: a definition that may have to be tied in to the density of floating virus particles, air circulation patterns, or whether or not contaminated surfaces are touched by susceptible individuals. In short, defining and measuring a contact or an effective contact turn out to be incredibly challenging (Mossong et al. 2008). That said, experimental methods may be used to estimate the average risk of acquiring, for example, tuberculosis (TB) or influenza, to individuals that spend on the average three hours per day in public transportation, in Mexico City or New York City.

In this paper, we propose the use of residence times in heterogeneous environments, as a proxy for “effective” contacts per unit time. Catching a communicable disease would of course depend on the presence of infected/infectious individuals (a necessary condition), the level of “risk” within a given “patch” (crowded bars, airports, schools, work places, etc.) and the time spent in such environment. Risk of infection is assumed to be a function of the time spent in pre-specified environments: risk that may be experimentally measured. We argue that characterizing a landscape as a collection of patches defined by risk (public transportation, schools, malls, work place, homes, etc.) is possible, especially if the risk of infection in such “local” environments is in addition a function of residence times and disease levels. Ranking patch-dependent risks of infection via the values of the transmission rate ( $\beta$ ) per unit of time may therefore be possible and useful. The reinterpretation of  $\beta$  and the use of residence times move us away from the world of models that account for transmission via the use of differential susceptibility to the world where infection depends on local environmental risk.

Consequently, we introduce a residence-time framework in the context of a multi-group system defined by patch-dependent risk (defined by  $\beta$ ). We study the role of patch residence times on disease dynamics within endemic and single-outbreak multi-group scenarios. Specifically, the study of the impact of patch residence times (modeled by a matrix of constants) on disease dynamics within a Susceptible-Infected-Susceptible (SIS) framework is carried out first, under the philosophy found in Brauer (2008), Brauer and Castillo-Chavez (1994), Brauer et al. (1996), Brauer and van den Driessche (2001), Brauer and Watmough (2009), Castillo-Chavez and Huang (1999), Haderler and Castillo-Chavez (1995) and Huang et al. (1992). Individuals move across patches as a function of their assessment of relative levels of infection in each area [studies using alternative classical approaches are found in Brauer and van den Driessche (2001), Brauer et al. (2008), Heiderich et al. (2002) and Velasco-Hernández et al. (1996)]. The concept of modeling disease dynamics where the population is structured into several communities goes back to Rushton and Mautner (1955). They considered an SI model with a constant population size in each community and derived solutions for their model. Multi-group models have surged in the literature to model sexually transmitted diseases. Lajmanovich and Yorke (1976) proposed an SIS model in the study of gonorrhea in a heterogeneous population. They obtained conditions to prove the global stability of both the disease-free equilibrium and the endemic equilibrium (EE). Nold (1980) proposed some extensions, allowing a more general contact form, of the model in Lajmanovich and Yorke (1976). Other multi-group models with different settings (including differential infectivity in each group) have been considered in Fall et al. (2007), Hethcote and Thieme (1985), Huang et al. (1992), Jacquez and Simon (1993), Jacquez et al. (1988, 1991), Lin and So (1993), Sattenspiel and Simon (1988) and Simon and Jacquez (1992). Hethcote and Thieme (1985) proved the uniqueness and the local stability of the EE if  $\mathcal{R}_0 > 1$  for an *SIRS* multi-group model. Lin and So (1993) proved the global stability of the EE if the effective contact rates between groups are small. Recently, authors in Fall et al. (2007) and Shuai and Driessche (2013) revisited the Lajmanovich and Yorke's model (Lajmanovich and Yorke 1976). Guo et al. (2006, 2008) used a combination of Lyapunov functions and elements of graph theory to prove the global stability of the EE of an SIR and SEIR multi-group models. Shuai and Driessche (2013) used a similar approach to study the asymptotic behavior of equilibria for some epidemic multi-group models. Typically, a sharp threshold property (Shuai and Driessche 2013), for which the disease dies out if  $\mathcal{R}_0 \leq 1$  and persists if  $\mathcal{R}_0 > 1$ , holds if the transmission (and contact) matrix  $B = (\beta_{ij})_{1 \leq i, j \leq n}$  is irreducible.

In this paper, we explore the disease dynamics when the residence-time matrix  $\mathbb{P}$  could be irreducible or not. First, we prove that the irreducibility of the residence-time matrix  $\mathbb{P}$  leads to a sharp threshold property. This property also holds when  $\mathbb{P}$  is replaced by the irreducible matrix  $\mathbb{P}D\mathbb{P}^t$  where  $D$  is a particular diagonal matrix and the residence-time matrix  $\mathbb{P}$  could be rectangular. Then, we study the disease dynamics at patch level by relaxing the irreducible condition on the residence-time matrix  $\mathbb{P}$ . Generalizations are explored through simulations of the two-patch SIS model with state-dependent residence times within our framework. The results are compared to the disease dynamics generated by constant residence times. More specifically,

the paper is organized as follows: Sect. 2 introduces a general  $n$  patch SIS model that accounts for residence times. Theoretical results on the role of residence time matrix ( $\mathbb{P}$ ) on disease dynamics are carried out using the residence-time-dependent basic reproduction number  $\mathcal{R}_0(\mathbb{P})$ . The patch-specific reproduction numbers  $\mathcal{R}_0^i(\mathbb{P})$ ,  $i = 1, \dots, n$  are defined to determine the disease persistence at the patch level. The usage of  $\mathcal{R}_0(\mathbb{P})$  allows us to explore the cases when the network configuration of patches is non-strongly connected. In addition, we also apply our framework to a SI model and a SIS model without demographics. Section 3 explores, through simulations, the dynamics of the SIS model under a state-dependent residence-time matrix in a two-patch system;  $\mathbb{P} \equiv \mathbb{P}(I_1, I_2)$ . That is, when the decisions to spend time in a patch are a function of patch-disease prevalence. Section 4 highlights our framework in the case of a two-patch single-outbreak SIR model following the work of Brauer (2008) and Brauer and Watmough (2009) and discusses the role of  $\mathbb{P}$  on the final epidemic size. Section 5 collects our observations, conclusions and discusses future work. The detailed proofs of our theoretical results are provided in the Appendix.

## 2 A General $n$ -Patch SIS Model with Residence Times

A general  $n$ -patch SIS model with residence-time matrix  $\mathbb{P}$  is derived. The global analysis of the model is carried out via the basic reproduction number  $\mathcal{R}_0$ . We also include patch-dependent disease persistence conditions.

### 2.1 Model Derivation

We model disease dynamics within an environment defined by  $n$  patches (or risk areas), and so, we let  $N_i(t)$ ,  $i = 1, 2, \dots, n$  denote resident population at Patch  $i$  at time  $t$ . We assume that Patch  $i$  residents spend  $p_{ij} \in [0, 1]$  time in Patch  $j$ , with  $\sum_{j=1}^{j=n} p_{ij} = 1$ , for each  $i = 1, \dots, n$ . In extreme cases, for example, we may have, for  $p_{ij} = 0$ ,  $i \neq j$ , that is Patch  $i$  residents spend no time in Patch  $j$  while  $\sum_{j \neq i} p_{ij} = 1$  (or equivalently  $p_{ii} = 0$ ) would imply that Patch  $i$  residents spend all their time in Patch  $j$  (with  $j = 1, \dots, n$  and  $j \neq i$ ) even though their patch is (labeled)  $i$ . In the absence of disease dynamics, the population of Patch  $i$  residents is modeled by the following equation:

$$\frac{dN_i}{dt} = b_i - d_i N_i \tag{1}$$

where the parameters  $b_i$  and  $d_i$  represent the birth rate and the natural per capita death rate in Patch  $i$ , respectively. Hence, the Patch  $i$  resident population approaches the constant  $\frac{b_i}{d_i}$  as  $t \rightarrow \infty$ .

In the presence of disease, we assume that disease dynamics are captured by an SIS model; thus, the Patch  $i$  resident population is divided into susceptible and infected classes, represented by  $S_i$ ,  $I_i$ , respectively, with  $S_i + I_i = N_i$ . We further assume that (a) there is no additional death due to disease; (b) the Patch  $i$  infected resident

population recovers and goes back to the susceptible class at the per capita rate  $\gamma_i$ ; (c) the residence-time matrix  $\mathbb{P} = (p_{ij})_{i=1, \dots, n}^{j=1, \dots, n}$  collects the proportion of times spent by  $i$ -residents in  $j$ -environments,  $i = 1, \dots, n$  and  $j = 1, \dots, n$ . The disease dynamics are therefore described by the following equations:

$$\begin{aligned} \dot{S}_i &= b_i - d_i S_i + \gamma_i I_i - \sum_{j=1}^n (S_i \text{ infected in Patch } j) \\ \dot{I}_i &= \sum_{j=1}^n (S_i \text{ infected in Patch } j) - \gamma_i I_i - d_i I_i \\ \dot{N}_i &= b_i - d_i N_i. \end{aligned} \tag{2}$$

We model  $S_i$  infection within Patch  $j$  in the following way:

- Since each  $p_{ij}$  entry of  $\mathbb{P}$  denotes the *proportion of time* that Patch  $i$  residents spent mingling in Patch  $j$ , we have that:
  - There are  $N_i p_{ij} = S_i p_{ij} + I_i p_{ij}$  Patch  $i$  residents in Patch  $j$  on the average at time  $t$ .
  - The total Patch  $j$ , the total effective population is  $\sum_{k=1}^n N_k p_{kj}$ , of which  $\sum_{k=1}^n I_k p_{kj}$  are infected. Hence, the proportion of infected individuals in Patch  $j$  is  $\frac{\sum_{k=1}^n I_k p_{kj}}{\sum_{k=1}^n N_k p_{kj}}$  and well defined, as long as there exists a  $k$  such that  $p_{kj} > 0$ , so that the population in Patch  $j$  is nonzero.
- Hence, the  $[S_i \text{ infected per unit of time in Patch } j]$  can be represented as the product of the following three items:

$$\begin{aligned} & \underbrace{\beta_j}_{\text{the risk of infection in Patch } j} \quad \times \quad \underbrace{S_i p_{ij}}_{\text{Susceptible from Patch } i \text{ who are currently in Patch } j} \\ & \times \quad \underbrace{\frac{\sum_{k=1}^n I_k p_{kj}}{\sum_{k=1}^n N_k p_{kj}}}_{\text{Proportion of infected in Patch } j}. \end{aligned}$$

The transmission takes on a modified frequency-dependent form that depends on how much time individuals of each epidemiological class spend in a particular area, where  $\beta_j$  differs by patch to reflect spatial differences in potential infectivity. More precisely,  $\beta_j$  is assumed to be a patch-specific measure of disease risk per unit of time with its effectiveness tied in to local environmental and sanitary conditions. Therefore,

$$[S_i \text{ infected per unit of time in Patch } j] \equiv \beta_j \times S_i p_{ij} \times \frac{\sum_{k=1}^n I_k p_{kj}}{\sum_{k=1}^n N_k p_{kj}} \tag{3}$$

provided that there exists  $k$  such that  $p_{kj} > 0$ .

Model (2) can be rewritten as follows:

$$\begin{aligned} \dot{S}_i &= b_i - d_i S_i + \gamma_i I_i - \sum_{j=1}^n \left( \beta_j \times S_i p_{ij} \times \frac{\sum_{k=1}^n I_k p_{kj}}{\sum_{k=1}^n N_k p_{kj}} \right), \\ \dot{I}_i &= \sum_{j=1}^n \left( \beta_j \times S_i p_{ij} \times \frac{\sum_{k=1}^n I_k p_{kj}}{\sum_{k=1}^n N_k p_{kj}} \right) - \gamma_i I_i - d_i I_i, \\ \dot{N}_i &= b_i - d_i N_i, \end{aligned} \tag{4}$$

with the dynamics of the Patch  $i$  resident total population modeled by the equation:  $\dot{N}_i(t) = b_i - d_i N_i(t)$ , where  $S_i + I_i = N_i$ , which implies that  $N_i(t) \rightarrow \frac{b_i}{d_i}$  as  $t \rightarrow +\infty$ . Theory of asymptotically autonomous systems for triangular systems (Castillo-Chavez and Thieme 1995; Vidyasagar 1980) guaranties that System (4) is asymptotically equivalent to:

$$\begin{aligned} \dot{i}_i &= \sum_{j=1}^n \left( \beta_j \left( \frac{b_i}{d_i} - I_i \right) p_{ij} \frac{\sum_{k=1}^n I_k p_{kj}}{\sum_{k=1}^n \frac{b_k}{d_k} p_{kj}} \right) - (\gamma_i + d_i) I_i \\ &= I_i \left( \frac{b_i}{d_i} - I_i \right) \left( \sum_{j=1}^n \frac{\beta_j p_{ij}^2}{\sum_{k=1}^n \frac{b_k}{d_k} p_{kj}} \right) \\ &\quad + \left( \frac{b_i}{d_i} - I_i \right) \sum_{j=1}^n \frac{\beta_j p_{ij} \sum_{k=1, k \neq i}^n I_k p_{kj}}{\sum_{k=1}^n \frac{b_k}{d_k} p_{kj}} - (d_i + \gamma_i) I_i \end{aligned} \tag{5}$$

for  $i = 1, 2, \dots, n$ , with residence-time matrix  $\mathbb{P} = (p_{ij})_{i=1, \dots, n}^{j=1, \dots, n}$  satisfying the conditions:

**HP1** At least one entry in each column of  $\mathbb{P}$  is strictly positive; and

**HP2** The sum of all entries in each row is one; i.e.,  $\sum_{j=1}^n p_{ij} = 1$  for all  $i$ .

**Remarks on Model (5):**

1. Timescales: We assume that the disease dynamics occurs at the comparable timescale as to the demographic dynamics and individuals enter or leave the patches at the relative faster timescale, e.g., daily or even hourly. The case when there is no demographics in the context of a single epidemic outbreak scenario, has been considered in Sect. 4.
2. In our current modeling framework, we assume that the residence-time matrix  $\mathbb{P}$  is a  $n \times n$  matrix. This approach could generalize the concept of  $k$  social groups and  $l$  patches by letting  $n = \max\{k, l\}$ ,  $p_{ij}|_{i>k} = 0$  and  $p_{ij}|_{j>l} = 0$ . The consequence of this generalization is that  $\mathbb{P}$  could have zero rows (when  $k < l = n$ ) or columns (when  $l < k = n$ ). The alternative treatment has been provided in the subsection 2.3 (thanks to the referee).
3. We do not assume that the residence-time matrix  $\mathbb{P}$  being irreducible, instead, we assume that it satisfies relaxed conditions **HP1–HP2**. More specifically, we will

explore the following two cases: (1) the global dynamics of Model (5) when the residence-time matrix  $\mathbb{P}$  is irreducible; (2) the persistence of disease dynamics at the patch level under conditions of **HP1** and **HP2** in the following subsection which includes the scenario when  $\mathbb{P}$  is not irreducible.

### 2.2 Equilibria, the Basic Reproduction Number and Global Analysis

To analyze the system, we investigate the basic reproduction number of the system with fixed residence times to better understand its properties in the absence of behavioral responses to risk. We let  $\mathcal{B} = (\beta_1, \beta_2, \dots, \beta_n)^t$  define the risk of infection vector;  $\beta_i$  is a measure of the risk per susceptible per unit of time while in residence in Patch  $i$ .

Letting  $S = (S_1, S_2, \dots, S_n)^t$ ,  $I = (I_1, I_2, \dots, I_n)^t$ ,  $\bar{N} = \left(\frac{b_1}{d_1}, \frac{b_2}{d_2}, \dots, \frac{b_n}{d_n}\right)^t$ ,

and  $\tilde{N} = \mathbb{P}^t \bar{N} = \begin{pmatrix} \sum_{k=1}^n \frac{b_k}{d_k} p_{k1} \\ \sum_{k=1}^n \frac{b_k}{d_k} p_{k2} \\ \vdots \\ \sum_{k=1}^n \frac{b_k}{d_k} p_{kn} \end{pmatrix}$ . Then, System (5) can be rewritten in the following compact (vectorial) form:

$$\dot{I} = \text{diag}(\bar{N} - I)\mathbb{P}\text{diag}(\mathcal{B})\text{diag}(\tilde{N})^{-1}\mathbb{P}^t I - \text{diag}(d_I + \gamma_I)I \tag{6}$$

with state space in  $\mathbb{R}_+^n$ . The set  $\Omega = \{I \geq \mathbf{0}_{\mathbb{R}^n}, I \leq \bar{N}\}$  is a compact positively invariant that attracts all trajectories of System (6). This implies that the populations involved are “biologically” well defined since solutions of (6) will converge to and stay in  $\Omega$ . We therefore restrict the dynamics of (6) to the compact set  $\Omega$ .

The analysis of System (6) is naturally tied in to the basic reproductive number  $\mathcal{R}_0$  (Diekmann et al. 1990; Driessche and Watmough 2002); the average number of secondary cases produced by an infected individual during its infectious period while interacting with a purely susceptible population.  $\mathcal{R}_0$  is given by (see the detailed formulation in Appendix):

$$\mathcal{R}_0 = \rho(-\text{diag}(\bar{N})\mathbb{P}\text{diag}(\mathcal{B})\text{diag}(\tilde{N})^{-1}\mathbb{P}^t V^{-1}) \tag{7}$$

where  $V = -\text{diag}(d_I + \gamma_I)$ ,  $d_I = (d_1, d_2, \dots, d_n)^t$  and  $\gamma_I = (\gamma_1, \gamma_2, \dots, \gamma_n)^t$ .

The basic reproduction number  $\mathcal{R}_0$  is used to establish global properties of System (6). For the relevant literature on global stability for multi-group or metapopulation models, see Arino and Driessche (2003), Iggidr et al. (2012), Kuniya and Muroya (2014), Lajmanovich and Yorke (1976), Sattenspiel and Simon (1988) and the references therein. We define the disease-free equilibrium (DFE) of System (6) as  $I^* = \mathbf{0}_{\mathbb{R}^n}$  and the endemic equilibrium (when  $\mathcal{R}_0 > 1$ ) as  $\bar{I}$  where all components are positive. By using the same approach as in Iggidr et al. (2012) and Lajmanovich and Yorke (1976), we arrive at the following theorem regarding the global dynamics of Model (6).

**Theorem 2.1** (Global dynamics of Model (6)) *Suppose that the residence-time matrix  $\mathbb{P}$  is irreducible, then the following statements hold:*

- If  $\mathcal{R}_0 \leq 1$ , the DFE  $I^* = \mathbf{0}_{\mathbb{R}^n}$  is globally asymptotically stable. If  $\mathcal{R}_0 > 1$  the DFE is unstable.
- If  $\mathcal{R}_0 > 1$ , there exists a unique endemic equilibrium  $\bar{I}$  which is GAS.

*Remarks* The detailed proof of Theorem 2.1 is provided in ‘‘Appendix 2.’’ These results imply that System (6) is robust; that is, disease outcomes are completely determined by whether or not the reproduction number  $\mathcal{R}_0$  is greater or less than one. The results of Theorem 2.1 while powerful do not provide easily accessible insights on the impact of the residence matrix  $\mathbb{P}$  on the levels of infection within each patch.

Direct insights on the effects of  $\mathbb{P}$  are derived by focusing on the levels of endemicity within each patch. The following two definitions help set the stage for the discussion:

- The basic reproduction number for Patch  $i$  in the absence of movement ( $p_{ii} = 1$  or  $\sum_{i \neq j} p_{ij} = 0$ ), SIS model, is defined as  $\mathcal{R}_0^i \equiv \frac{\beta_i}{d_i + \gamma_i}$ , which determines whether or not the disease will be endemic in Patch  $i$ . In short disease will die out if  $\mathcal{R}_0^i \leq 1$  with a unique endemic equilibrium, that is GAS, if  $\mathcal{R}_0^i > 1$ .
- The basic reproduction number associated with Patch  $i$ , under the presence of multi-patch residents, is defined as follows:

$$\begin{aligned} \mathcal{R}_0^i(\mathbb{P}) &= \frac{\sum_{j=1}^n \beta_j \left(\frac{b_i}{d_i} p_{ij}\right) \left(\frac{p_{ij}}{\sum_{k=1}^n \frac{b_k}{d_k} p_{kj}}\right)}{d_i + \gamma_i} = \frac{\sum_{j=1}^n p_{ij} \beta_j \left(\frac{\left(\frac{b_i}{d_i} p_{ij}\right)}{\sum_{k=1}^n \frac{b_k}{d_k} p_{kj}}\right)}{d_i + \gamma_i} \\ &= \mathcal{R}_0^i \times \sum_{j=1}^n p_{ij} \left(\frac{\beta_j}{\beta_i}\right) \left(\frac{\left(\frac{b_i}{d_i} p_{ij}\right)}{\sum_{k=1}^n \frac{b_k}{d_k} p_{kj}}\right). \end{aligned}$$

We explore the role that  $\mathcal{R}_0^i(\mathbb{P})$  plays in determining the impact of all residents on disease dynamics persistence in Patch  $i$  in the following theorem.

**Theorem 2.2** (The endemicity of disease in Patch  $i$ ) *Assume that the residence-time matrix  $\mathbb{P}$  satisfies Condition **HP1** and **HP2** but that some of its entries can be zeros.*

- If  $\mathcal{R}_0^i(\mathbb{P}) > 1$ , then the disease persists in Patch  $i$ .
- If the following conditions hold:

$$\mathbf{H}: p_{kj} = 0 \text{ for all } k = 1, \dots, n, \text{ and } k \neq i, \text{ whenever } p_{ij} > 0,$$

then we have

$$\mathcal{R}_0^i(\mathbb{P}) = \mathcal{R}_0^i \times \sum_{j=1}^n p_{ij} \left(\frac{\beta_j}{\beta_i}\right) \left(\frac{\left(p_{ij} \frac{b_i}{d_i}\right)}{\sum_{k=1}^n \frac{b_k}{d_k} p_{kj}}\right) = \mathcal{R}_0^i \times \sum_{j=1}^n p_{ij} \left(\frac{\beta_j}{\beta_i}\right).$$

Thus, when Condition **H** holds and  $\mathcal{R}_0^i \times \sum_{j=1}^n p_{ij} \left(\frac{\beta_j}{\beta_i}\right) < 1$ , then endemic levels of disease cannot be supported in Patch  $i$ . That is,

$$\lim_{t \rightarrow \infty} I_i(t) = 0.$$

*Remarks* The detailed proof of Theorem 2.2 is provided in ‘‘Appendix 3.’’ The results of Theorem 2.2 give insights on the role that the infection risk (measured by  $\mathcal{B}$ ) and the residence-time matrix ( $\mathbb{P}$ ) have in promoting or suppressing infection. Further, a closer look at the expression of the general basic reproduction number in Patch  $i$ , namely

$$\mathcal{R}_0^i(\mathbb{P}) = \mathcal{R}_0^i \times \sum_{j=1}^n p_{ij} \left(\frac{\beta_j}{\beta_i}\right) \left(\frac{\left(\frac{b_i}{d_i} p_{ij}\right)}{\sum_{k=1}^n \frac{b_k}{d_k} p_{kj}}\right),$$

leads to the following observations:

1. The movement between patches, modeled via residence-time matrix  $\mathbb{P}$ , can promote endemicity: For example, if  $\mathcal{R}_0^i = \frac{\beta_i}{d_i + \gamma_i} \leq 1$ , i.e., there is no endemic disease in Patch  $i$ , then the presence of movement connecting Patch  $i$  to possibly all other patches can support endemic disease levels in the following ways:
  - Via the presence of high-risk patches, that is, there exists a patch  $j$  such that  $\frac{\beta_j}{\beta_i}$  is large enough. For example, letting  $p_{kl} = 1/n$  for all  $k, l$  with the total population in each patch being the same ( $\frac{b_k}{d_k} = K$  for all  $k$ ;  $K$  a constant) then  $\mathcal{R}_0^i(\mathbb{P}) = \mathcal{R}_0^i \frac{\sum_{j=1}^n \beta_j}{n\beta_i}$ , and consequently, if  $\sum_{j=1}^n \beta_j > \frac{n\beta_i}{\mathcal{R}_0^i}$ , then Patch  $i$  will promote the disease at endemic levels.
  - Whenever individuals spend more time in high-risk patches than in low-risk patches. For example, in the extreme case,  $p_{ij} = 1$  with  $\frac{\beta_j}{\beta_i} > \frac{1}{\mathcal{R}_0^i}$ , we have that  $\mathcal{R}_0^i(\mathbb{P}) > 1$ , and thus, endemic disease levels in Patch  $i$  can be supported. Patch  $j$  ( $j = 1, \dots, n$  and  $j \neq i$ ) can therefore be considered the source and Patch  $i$  ( $i \neq j$ ) the sink (Arino 2008; Arino et al. 2005; Arino and Driessche 2003, 2006; Kuniya and Muroya 2014; Lajmanovich and Yorke 1976; Sattenspiel and Simon 1988; Sattenspiel and Dietz 1995).
2. Under the assumption  $\mathcal{R}_0^i > 1$ , for an isolated Patch  $i$ , conditions that lead to disease extinction in the same Patch  $i$  under the movement can be identified. According to Theorem 2.2, Condition **H** should be satisfied and so the expression of  $\mathcal{R}_0^i(\mathbb{P})$  reduces to

$$\mathcal{R}_0^i(\mathbb{P}) = \mathcal{R}_0^i \times \sum_{j=1}^n p_{ij} \left(\frac{\beta_j}{\beta_i}\right) \left(\frac{\left(\frac{b_i}{d_i} p_{ij}\right)}{\sum_{k=1}^n \frac{b_k}{d_k} p_{kj}}\right) = \mathcal{R}_0^i \times \sum_{j=1}^n p_{ij} \left(\frac{\beta_j}{\beta_i}\right).$$

Therefore, the only way to have the value of  $\mathcal{R}_0^i(\mathbb{P})$  be less than one, would be when the amount of time spent in Patch  $i$  is such that  $\sum_{j=1}^n \left(\frac{\beta_j}{\beta_i}\right) p_{ij} < \frac{1}{\mathcal{R}_0^i(\mathbb{P})}$ . Therefore, we conclude that the synergy between the residence-time matrix  $\mathbb{P}$  and the existence of sufficient low-risk patches (i.e.,  $\beta_j \ll \beta_i$ ) can suppress a disease outbreak in Patch  $i$ .

### 2.3 Social Groups Versus Patch Environments

We assume that there are  $n$  social groups interacting in  $m$  different patch environments. Let  $p_{ij}$  be the proportion time of social group  $i$  spent at patch environment  $j$ , then the residence-time matrix  $\mathbb{P} = (p_{ij})_{\substack{1 \leq i \leq n \\ 1 \leq j \leq m}}$  is a  $n \times m$  matrix.

Following the same modeling approach of the system (5), Model 5 is rewritten as the following form:

$$\begin{aligned} \dot{I}_i &= \left(\frac{b_k}{d_k} - I_i\right) \sum_{j=1}^m \beta_j p_{ij} \frac{\sum_{k=1}^n I_k p_{kj}}{\sum_{l=1}^n \frac{b_l}{d_l} p_{lj}} - (d_i + \gamma_i) I_i \\ &= \left(\frac{b_k}{d_k} - I_i\right) \sum_{k=1}^n \left(\frac{\sum_{j=1}^m p_{ij} \beta_j p_{kj}}{\sum_{l=1}^n \frac{b_l}{d_l} p_{lj}}\right) I_k - (d_i + \gamma_i) I_i \\ &= \left(\frac{b_k}{d_k} - I_i\right) \sum_{k=1}^n b_{ik} I_k - (d_i + \gamma_i) I_i \end{aligned} \tag{8}$$

where

$$b_{ik} = \sum_{j=1}^m \left(\frac{p_{ij} \beta_j p_{kj}}{\sum_{l=1}^n \frac{b_l}{d_l} p_{lj}}\right).$$

Model (8) is isomorphic to those considered in Fall et al. (2007), Guo et al. (2006), Guo et al. (2008) and Shuai and Driessche (2013), which could be rewritten in the simplified form (6) as Model (5). Denote the disease transmission matrix  $B = (b_{ik})_{1 \leq i, k \leq n}$ , then we also have the form of  $B = \mathbb{P} \text{diag}(\mathcal{B}) \text{diag}(\tilde{N})^{-1} \mathbb{P}^t$  which is symmetric. We could see that Theorem 2.1 still holds if the irreducibility of  $\mathbb{P}$  is replaced by the irreducibility of  $B$ . We should also expect similar results of Theorem 2.2 for Model (8) when  $B$  is not irreducible.

The difference between the models in the aforementioned papers (e.g., [Fall et al. 2007](#); [Guo et al. 2006, 2008](#); [Shuai and Driessche 2013](#)) and our model (8) is that: In the former models, the disease transmission coefficient  $\beta_{ij}$  involves the contact rate between group  $j$  and group  $i$ ; see the case of bilinear incidence ([Guo et al. 2006, 2008](#); [Shuai and Driessche 2013](#)) and proportional of  $\beta_{ij}$  for the frequency-dependent incidence ([Fall et al. 2007](#); [Guo et al. 2006](#)). In our case, the disease transmission matrix  $B = (b_{ik})_{1 \leq i, k \leq n}$  is symmetric and incorporates the environmental risks in different patches and the proportion of times that different social groups spent in each patch.

Now we apply the approach above to a SIS model without demographics and a SI model with the disease-induced death rate  $c_i$  for each social group  $i$  as follows.

1. A SIS model without demographics (i.e., assume that the total population size  $N_i$  at each patch  $i$  is constant and the natural death rate  $d_i = 0$ ) could be rewritten in the form of:

$$\dot{I}_i = S_i \sum_{k=1}^n b_{ik} I_k - \gamma_i I_i, \quad S_i = N_i - I_i \tag{9}$$

whose simplified form is

$$\dot{I} = SBI - GI$$

where  $B = (b_{ik})_{1 \leq i, k \leq n} = \left( \sum_{j=1}^m \left( \frac{p_{ij} \beta_j p_{kj}}{n} \right) \right)_{1 \leq i, k \leq n}$  is symmetric, and

$G = (\gamma_i \delta_{ik})_{1 \leq i, k \leq n}$  with  $\delta_{ik}$  being the Kronecker delta. The basic reproduction number of Model (9) is  $\rho(BG^{-1})$  which determines whether the disease persists or dies out. Both our Theorems 2.1 and 2.2 could be applied to Model (9)

2. A SI model with the disease-induced death rate  $c_i$  for each social group  $i$  could be rewritten in the form of:

$$\dot{I}_i = S_i \sum_{k=1}^n b_{ik} I_k - c_i I_i, \quad S_i = N_i - I_i \tag{10}$$

whose simplified form is

$$\dot{I} = SBI - CI$$

where  $C = (c_i \delta_{ik})_{1 \leq i, k \leq n}$ , and  $B = (b_{ik})_{1 \leq i, k \leq n} = \left( \sum_{j=1}^m \left( \frac{p_{ij} \beta_j p_{kj}}{n} \right) \right)_{1 \leq i, k \leq n}$  is still symmetric but depending on the total population size  $N_i$  in each patch.

Notice that  $\dot{N}_l = -c_l I_l$ , we expect the total population at each patch approaches zero as time is large enough. This is confirmed by simulations. Simulations also suggest that the infection dynamics have similar patterns as the prevalence of the SIR model studied in Sect. 4, and the limit of  $I_i(t)/N_i(t)$  goes to 1 for each patch as time is large enough.

### 3 Two-Patch Models: State-Dependent Residence-Time Matrix

We now extend the analysis of disease dynamics to the case where susceptible individuals respond to variations in risk in an automatic way. In particular, we consider the case when susceptible individuals make programmed responses to variations in disease risk, and do not choose their response to optimize an index of well-being (see, e.g., Brauer 2008; Brauer et al. 1996; Brauer and van den Driessche 2001; Brauer and Watmough 2009). While this may not be a very good approximation of disease risk management in real systems, it enables us to explore the implications of certain types of phenomenologically modeled behavioral responses by assuming, for example, that the proportion of time spent in a particular patch depends on the numbers of infected individuals on that particular patch; that is  $\mathbb{P} \equiv \mathbb{P}(I_1, I_2)$ .

Possible properties of the proportion of time spent by resident of Patch  $i$  into Patch  $j$ ,  $i \neq j$ ,  $(p_{ij})$  may include: increases with respect to the growth of infected resident in Patch  $i$  ( $I_i$ ) or decreases with respect to infected resident in Patch  $j$  ( $I_j$ ). Mathematically, we would have that

$$\frac{\partial p_{ij}(I_i, I_j)}{\partial I_j} \leq 0 \quad \text{and} \quad \frac{\partial p_{ij}(I_i, I_j)}{\partial I_i} \geq 0.$$

In a two-patch system, the use of the relationship  $p_{ij}(I_1, I_2) + p_{ji}(I_1, I_2) = 1$  reduces the above four conditions on  $\mathbb{P}$  to the following conditions:

$$\frac{\partial p_{11}(I_1, I_2)}{\partial I_1} \leq 0 \quad \text{and} \quad \frac{\partial p_{22}(I_1, I_2)}{\partial I_2} \leq 0.$$

Examples of functions  $p_{ij}(I_1, I_2)$  with these properties include,

$$p_{12}(I_1, I_2) = \sigma_{12} \frac{1 + I_1}{1 + I_1 + I_2} \quad \text{and} \quad p_{21}(I_1, I_2) = \sigma_{21} \frac{1 + I_2}{1 + I_1 + I_2}$$

and

$$p_{11}(I_1, I_2) = \frac{\sigma_{11} + \sigma_{11}I_1 + I_2}{1 + I_1 + I_2} \quad \text{and} \quad p_{22}(I_1, I_2) = \frac{\sigma_{22} + I_1 + \sigma_{22}I_2}{1 + I_1 + I_2}$$

where  $\sigma_{ij}$  are such that  $\sum_{j=1}^2 \sigma_{ij} = 1$ .

More complex behavioral response formulations may also depend on the states of total populations  $N_1$  and  $N_2$ , but the current specification captures important components of risk (infections) and allows us to retain the asymptotic equivalence property applied in the case of fixed residence times. Hence, using the same notation as in System (6) leads to the following two-dimensional system with  $\mathbb{P} = \mathbb{P}(I_1, I_2)$ :

$$\begin{cases} \dot{I}_1 = X(I_1, I_2) \left(\frac{b_1}{d_1} - I_1\right) I_1 + Y(I_1, I_2) \left(\frac{b_1}{d_1} - I_1\right) I_2 - (d_1 + \gamma_1) I_1, \\ \dot{I}_2 = Y(I_1, I_2) \left(\frac{b_2}{d_2} - I_2\right) I_1 + Z(I_1, I_2) \left(\frac{b_2}{d_2} - I_2\right) I_2 - (d_2 + \gamma_2) I_2, \end{cases} \tag{11}$$

where

$$\begin{aligned} X(I_1, I_2) &= \frac{\beta_1 p_{11}^2(I_1, I_2)}{p_{11}(I_1, I_2) \frac{b_1}{d_1} + p_{21}(I_1, I_2) \frac{b_2}{d_2}} + \frac{\beta_2 p_{12}^2(I_1, I_2)}{p_{12}(I_1, I_2) \frac{b_1}{d_1} + p_{22}(I_1, I_2) \frac{b_2}{d_2}}, \\ Y(I_1, I_2) &= \frac{\beta_1 p_{11}(I_1, I_2) p_{21}(I_1, I_2)}{p_{11}(I_1, I_2) \frac{b_1}{d_1} + p_{21}(I_1, I_2) \frac{b_2}{d_2}} + \frac{\beta_2 p_{12}(I_1, I_2) p_{22}(I_1, I_2)}{p_{12}(I_1, I_2) \frac{b_1}{d_1} + p_{22}(I_1, I_2) \frac{b_2}{d_2}}, \end{aligned}$$

and

$$Z(I_1, I_2) = \frac{\beta_1 p_{21}^2(I_1, I_2)}{p_{11}(I_1, I_2) \frac{b_1}{d_1} + p_{21}(I_1, I_2) \frac{b_2}{d_2}} + \frac{\beta_2 p_{22}^2(I_1, I_2)}{p_{12}(I_1, I_2) \frac{b_1}{d_1} + p_{22}(I_1, I_2) \frac{b_2}{d_2}},$$

where  $X(I_1, I_2)$ ,  $Y(I_1, I_2)$  and  $Z(I_1, I_2)$  are positive functions of  $I_1$  and  $I_2$ .

The basic reproduction number  $\mathcal{R}_0$  is the same as in the previous section since it is computed at the infection-free state, i.e.,

$$\mathcal{R}_0 = \rho(\text{diag}(\tilde{N}) \mathbb{P} \text{diag}(\mathcal{B}) \text{diag}(\tilde{N})^{-1} \mathbb{P}^t (-V^{-1}))$$

where, in this case, we have that  $\mathbb{P} = \begin{bmatrix} \sigma_{11} & \sigma_{12} \\ \sigma_{21} & \sigma_{22} \end{bmatrix}$  and  $\sigma_{ij} = p_{ij}(0, 0)$ ,  $\forall \{i, j\} = \{1, 2\}$ .

The properties of positiveness and boundedness of trajectories of System (6) are preserved in System (11). In addition, System (11) has a unique DFE equilibrium whose local stability is determined by the value of the  $\mathcal{R}_0$ : The DFE is locally asymptotically stable if  $\mathcal{R}_0 < 1$  while it is unstable if  $\mathcal{R}_0 > 1$ .

Let us consider whether System (11) can have a boundary equilibrium such as  $(0, \bar{I}_2)$  or  $(\bar{I}_1, 0)$ . The assumption that System (11) has such a boundary equilibrium  $(0, \bar{I}_2)$  with  $\bar{I}_2 > 0$  implies that  $Y(0, \bar{I}_2) = 0$ . Since  $p_{11}(0, I_2) = \frac{\sigma_{11} + I_2}{1 + I_2}$  and  $p_{22}(0, I_2) = \sigma_{22}$ , we deduce that

$$Y(0, I_2) = \frac{\beta_1 \sigma_{21} (\sigma_{11} + I_2)}{\frac{\sigma_{11} + I_2}{1 + I_2} \frac{b_1}{d_1} + \sigma_{21} \frac{b_2}{d_2} (1 + I_2)} + \frac{\beta_2 \sigma_{12} \sigma_{22}}{\sigma_{12} \frac{b_1}{d_1} + \sigma_{22} \frac{b_2}{d_2} (1 + I_2)}.$$

This indicates that  $Y(0, I_2) = 0$  if and only if  $\sigma_{21} = 0$  and  $\sigma_{12} = 0$ , which requires that:

$$p_{12} = p_{21} = 0, \text{ and } p_{11} = p_{22} = 1.$$

A similar arguments can be applied to the boundary equilibrium  $(\bar{I}_1, 0)$ . Therefore, we conclude that System (11) will have a boundary equilibrium  $(0, \bar{I}_2)$  or  $(\bar{I}_1, 0)$  only in the trivial case of isolated patches, that is, where there is no movement between two patches. This conclusion differs from the state-independent residence matrix model (6), since for example, the two-patch model (6), according to Theorem 2.1, boundary equilibrium  $(0, \bar{I}_2)$  or  $(\bar{I}_1, 0)$  can exist when  $p_{11} = p_{22} = 0$  ( $p_{12} = p_{21} = 1$ ).

To illustrate the difference between the state-dependent residence matrix model (11) and the state-independent residence matrix model (6), we look at the situation when  $\sigma_{11} = \sigma_{22} = 0, \sigma_{12} = \sigma_{21} = 1$  ( $p_{11} = p_{22} = 0, p_{12} = p_{21} = 0$  for the state-independent residence matrix model (6)). Under the condition of  $\sigma_{11} = \sigma_{22} = 0, \sigma_{12} = \sigma_{21} = 1$ , we have Model (11), that

$$p_{12}(I_1, I_2) = \frac{1 + I_1}{1 + I_1 + I_2} \text{ and } p_{21}(I_1, I_2) = \frac{1 + I_2}{1 + I_1 + I_2}$$

and


$$p_{11}(I_1, I_2) = \frac{I_2}{1 + I_1 + I_2} \text{ and } p_{22}(I_1, I_2) = \frac{I_1}{1 + I_1 + I_2}.$$

This difference has significant impact on disease dynamics (see Fig. 1a, b, red curves).


In Fig. 1b, we see that the infection in Patch 2 (high risk) persists in the state-dependent case, whereas it dies out when  $\mathbb{P}$  is constant. That is due to the fact that  $p_{ii}(I_1, I_2)$  will not equal zero, whereas  $p_{ij}(I_1, I_2)$  with  $i \neq j$  may. For the constant residence-time matrix, the dynamics of the disease in each patch is also independent, where people in Patch  $i$  infect only susceptible in Patch  $j$  with  $i \neq j$ . In Fig. 1b (red solid curve), we observe that the disease dies out in Patch 2 with  $\tilde{\mathcal{R}}_0^2 = \frac{\beta_1}{d_2 + \gamma_2} = 0.8571$ . For the state-dependent case, unless there is no disease in both patches or one disease-free patch, the proportion of time residents spend in their own patch is nonzero. This leads the disease to persist in both patches if  $\mathcal{R}_0 > 1$  (see Fig. 1b, red dashed curves). However, even in this case, the disease dies out in both patches if  $\mathcal{R}_0 < 1$  (see Fig. 3, red curves, for instance).

### 3.1 Applications and Comparisons: The Two-Patch Cases

The analytical results of the global dynamics on the asymptotic behavior of Model (11) are still unresolved. Hence, we ran simulations to gain some insights on the role of  $\mathbb{P}(I_1, I_2)$  on endemic dynamics. We observe that trajectories converge toward an endemic equilibrium whenever  $\mathcal{R}_0 > 1$ ; however, there are substantial differences in the transient dynamics generated by state-dependent  $\mathbb{P}(I_1, I_2)$  when compared to those generated with a constant residence-time matrix.




**Fig. 1** Coupled dynamics of  $I_1$  and  $I_2$  for constant  $p_{ij}$  (solid) and state-dependent  $p_{ij}$  (dashed). The red lines is case of high mobility, i.e.,  $p_{12} = p_{21} = \sigma_{12} = \sigma_{21} = 1$ . The black lines represent the symmetric case, i.e.,  $p_{12} = \sigma_{12} = 0.5$ , and the blue line represent the polar case, i.e.,  $p_{12} = p_{21} = \sigma_{12} = \sigma_{21} = 0$ . **a** Dynamics of the disease in Patch 1. If there is no movement between the patches (blue curves), the disease dies out in the low-risk Patch 1 in both approaches with  $\mathcal{R}_0^1 = 0.7636$ . **b** Dynamics of the disease in Patch 2. In the high-mobility case, the disease dies out (solid red curve) for  $\mathbb{P}$  constant, with  $\mathcal{R}_0^2 = 0.8571$ , and persists for  $\mathbb{P}$  state-dependent (dashed red curve) (Color figure online)



**Fig. 2** Coupled dynamics of  $I_1 + I_2$  for constant  $p_{ij}$  (solid) and state-dependent  $p_{ij}$  (dashed). The overall prevalence is higher if the residence times is symmetric (solid and dashed black curves). The black curves represent the symmetric case ( $p_{12} = p_{21} = \sigma_{12} = \sigma_{21} = 0.5$ ), and the blue lines represent the polar case ( $p_{12} = p_{21} = \sigma_{12} = \sigma_{21} = 0$ ) and red curves represent high-mobility case ( $p_{12} = p_{21} = \sigma_{12} = \sigma_{21} = 1$ ) (Color figure online)

Unless stated otherwise, we suppose the following generic values for the simulations:  $\beta_1 = 0.3$ ,  $\beta_2 = 1.2$ ,  $b_1 = 9$ ,  $d_1 = 1/7$ ,  $b_2 = 9$ ,  $d_2 = 1/10$  and  $\gamma_1 = \gamma_2 = 1/4$ . We carried out numerical simulation for a range of residence-time matrices. It is observed that:

1. For the symmetric case where  $p_{12} = p_{21} = 0.5$ , the disease is endemic in both patches as predicted by Theorem 2.1 since  $\mathcal{R}_0 = 2.0466$ . For the state-dependent case, simulations suggest (Fig. 1a, b, black dashed curves) that trajectories tend to be endemic in both patches. However, the level of endemicity is lower than the constant case in Patch 1 (low-risk patch) and is greater in Patch 2 (high-risk patch).
2. Figure 2 sketches the overall prevalence in both patches with three different scenarios of residence-time matrix  $\mathbb{P}$ , both the constant and state-dependent case. The disease persists since the overall  $\mathcal{R}_0 > 1$  in all three cases.
3. The case where there is no movement between patches, that is,  $p_{12} = p_{21} = 0$  ( $p_{11} = p_{22} = 1$ ) and  $\sigma_{12} = \sigma_{21} = 0$  (or  $p_{12}(I_1, I_2) = p_{21}(I_1, I_2) = 0$ ), corresponds to the case where the system behaves as two isolated patches. In this case, the disease dies out or persists in Patch  $i$  if  $\mathcal{R}_0^i$  is above or below unity in both approaches. This is illustrated in Fig. 1a, b) where the disease dies out in Patch 1 ( Fig. 1a, blue solid line) where  $\mathcal{R}_0^1 = \frac{\beta_1}{d_1 + \gamma_1} = 0.7636$  and the disease persists in Patch 2 (Fig. 1b, blue solid curve) where  $\mathcal{R}_0^2 = \frac{\beta_2}{d_2 + \gamma_2} = 3.4286$ . For the state-dependent case (dashed blue curves in Fig. 1a, b), the outcome is similar to the constant residence-time case.
4. In Fig. 4a, b, we explore the cases where there is symmetry ( $\sigma_{ij} = \sigma_{ji}$ ) with  $\sigma_{ij} = p_{ij}(0, 0)$ . We supposed in this case that Patch 2 has higher risk ( $\beta_2 = 1.2$ ) and Patch 1 has lower risk ( $\beta_1 = 0.3$ ). As can be intuitively deduced, the prevalence in Patch 1 is at its highest in the case of “high mobility” ( $\sigma_{12} = \sigma_{21} = 1$ ), and decreasing as  $\sigma_{ij}$  decreases (with  $i \neq j$ ). Conversely, prevalence in Patch 2 is at its highest under very “low mobility” ( $\sigma_{12} = \sigma_{21} = 0$ ) and decreases as  $\sigma_{ij}$




**Fig. 3** Dynamics of  $I_1$  and  $I_2$  for varying  $\sigma_{ij}$  for the state-dependent  $p_{ij}(I_1, I_2)$  where  $\mathcal{R}_0 < 1$ . This is obtained by using the values  $\beta_1 = 0.2$  and  $\beta_2 = 0.3$ . In all the three cases, the disease dies out in both patches. The *black curves* represent the symmetric case ( $p_{12} = p_{21} = \sigma_{12} = \sigma_{21} = 0.5$ ), the *blue line* represent the polar case ( $p_{12} = p_{21} = \sigma_{12} = \sigma_{21} = 0$ ), and *red curves* represent high-mobility case ( $p_{12} = p_{21} = \sigma_{12} = \sigma_{21} = 1$ ) (Color figure online)

increases. Note that  $\sigma_{ij}$ , with  $i \neq j$  is proportional to  $p_{ij}(I_1, I_2)$  which is the actual residence time.


- We continue to explore the asymmetric case ( $\sigma_{ij} \neq \sigma_{ji}$ ); that is, there is more mobility toward one patch. In Fig. 5a, the prevalence in Patch 1 (low risk) is at its highest if there is “high mobility” from Patch 1 to Patch 2 ( $\sigma_{12} = 1$ ) and no mobility from Patch 2 to Patch 1 ( $\sigma_{21} = 0$ ), and the prevalence decreases along with  $\sigma_{12}$ . If the programmed response of residents of Patch 1 is to reduce their mobility ( $\sigma_{12} = 0$ ) then, even if the mobility of residents in the high-risk Patch 2 is extremely high ( $\sigma_{21} = 1$ ), still the prevalence in Patch 1 is at its lowest. Similar remarks hold for Fig. 5b regarding the prevalence in Patch 2 (high risk) under different mobility schemes.
- Finally, Fig. 6 presents the dynamics of the infected in both patches for the (conventional) case where  $p_{12} = 0$  (and  $p_{11} = 1$ ). This case is particularly interesting since the residence-time matrix  $\mathbb{P}$  is not irreducible (hence, the hypothesis of Theorem 2.1 fails) but  $\mathcal{R}_0^i(\mathbb{P}) = 1.8929 > 1$ . As predicted by Theorem 2.2, the disease in Patch 2 is persistent. Also, it is worth noticing that in Fig. 6,  $I_1$  persists as well even though  $\mathcal{R}_0^1(\mathbb{P}) = 0.4455 < 1$ , as the condition  $\mathcal{R}_0^i(\mathbb{P}) > 1$ , for  $i = 1, 2$ , is sufficient but not necessary for persistence in Patch  $i$ .

## 4 Final Epidemic Size


The study of the role of residence-time matrices on the dynamics of a single outbreak within a Susceptible-Infected-Recovered (with immunity) or SIR model without births and deaths is relevant to the development of public disease management measures (Brauer et al. 2010; Chowell et al. 2015; Hernandez-Ceron et al. 2013). Under the parameters and definitions introduced earlier, and making use of the same notation, we arrive at the following system of nonlinear differential equations:



**Fig. 4** Dynamics of  $I_1$  and  $I_2$  for varying  $\sigma_{ij}$  for the state-dependent  $P_{ij}(I_1, I_2)$  approach. **a** The level of prevalence in Patch 1 (low risk) seems to decrease as  $\sigma_{12}$  and  $\sigma_{21}$  decrease. **b** The level of prevalence in Patch 2 (high risk) seems to increase as  $\sigma_{12}$  and  $\sigma_{21}$  decrease (Color figure online)



**Fig. 5** Dynamics of  $I_1$  and  $I_2$  for varying  $\sigma_{ij}$ , but non-symmetric, for the state-dependent  $p_{ij}(I_1, I_2)$ . **a** The level of prevalence in Patch 1 (low risk) seems to decrease as  $\sigma_{12}$  and  $\sigma_{21}$  decrease. **b** The level of prevalence in Patch 2 (high risk) seems to increase as  $\sigma_{12}$  and  $\sigma_{21}$  decrease (Color figure online)



**Fig. 6** Dynamics of  $I_1$  and  $I_2$  where  $p_{12} = 0$ . In this case, the residence-time matrix  $\mathbb{P}$  is not irreducible, the disease in Patch 2 persists nonetheless as predicted by the Theorem 2.2 (Color figure online)

$$\begin{cases} \dot{S}_i = - \left( \frac{\beta_i p_{ii}^2}{N_i p_{ii} + N_j p_{ji}} + \frac{\beta_j p_{ij}^2}{N_i p_{ij} + N_j p_{jj}} \right) S_i I_i - \left( \frac{\beta_i p_{ii} p_{ji}}{N_i p_{ii} + N_j p_{ji}} + \frac{\beta_j p_{ij} p_{jj}}{N_i p_{ij} + N_j p_{jj}} \right) S_i I_j, \\ \dot{I}_i = \left( \frac{\beta_i p_{ii}^2}{N_i p_{ii} + N_j p_{ji}} + \frac{\beta_j p_{ij}^2}{N_i p_{ij} + N_j p_{jj}} \right) S_i I_i + \left( \frac{\beta_i p_{ii} p_{ji}}{N_i p_{ii} + N_j p_{ji}} + \frac{\beta_j p_{ij} p_{jj}}{N_i p_{ij} + N_j p_{jj}} \right) S_i I_j - \alpha_i I_i, \\ \dot{R}_i = \alpha_i I_i, \end{cases} \tag{12}$$


where  $R_i$  denotes the population of recovered immune individuals in Patch  $i$ ,  $\alpha_i$  is the recovery rate in Patch  $i$ , and  $N_i \equiv S_i + I_i + R_i$ , for  $i = 1, 2$ .

The basic reproduction number  $\mathcal{R}_0$  is by definition the largest eigenvalue of  $2 \times 2$  ( $n \times n$  for the general case) next-generation matrix,

$$-FV^{-1} = \begin{pmatrix} \left( \frac{\beta_1 p_{11}^2}{N_1 p_{11} + N_2 p_{21}} + \frac{\beta_2 p_{12}^2}{N_1 p_{12} + N_2 p_{22}} \right) \frac{N_1}{\alpha_1} & \left( \frac{\beta_1 p_{11} p_{21}}{N_1 p_{11} + N_2 p_{21}} + \frac{\beta_2 p_{12} p_{22}}{N_1 p_{12} + N_2 p_{22}} \right) \frac{N_1}{\alpha_2} \\ \left( \frac{\beta_1 p_{11} p_{21}}{N_1 p_{11} + N_2 p_{21}} + \frac{\beta_2 p_{12} p_{22}}{N_1 p_{12} + N_2 p_{22}} \right) \frac{N_2}{\alpha_1} & \left( \frac{\beta_1 p_{21}^2}{N_1 p_{11} + N_2 p_{21}} + \frac{\beta_2 p_{22}^2}{N_1 p_{12} + N_2 p_{22}} \right) \frac{N_2}{\alpha_2} \end{pmatrix}.$$

It has been shown (see [Hethcote 1976](#), for example) that not everybody gets infected during an outbreak, and so, estimating the size of the recovered population (the final epidemic size in the absence of deaths or departures) is tied in the solutions of the final size relationship, given in this case, by the system:

$$\begin{bmatrix} \log \frac{S_1(0)}{S_1(\infty)} \\ \log \frac{S_2(0)}{S_2(\infty)} \end{bmatrix} = \begin{bmatrix} K_{11} & K_{12} \\ K_{21} & K_{22} \end{bmatrix} \begin{bmatrix} 1 - \frac{S_1(\infty)}{N_1} \\ 1 - \frac{S_2(\infty)}{N_2} \end{bmatrix} \tag{13}$$



**Fig. 7** Prevalence in patch 1 (low risk) reaches its highest when in extreme mobility case (*solid blue line*) and is lowest when there is no mobility between the patches. The opposite of this scenario happens in patch 2 (high risk) (*Color figure online*)

where

$$K = \begin{bmatrix} \left( \frac{\beta_1 p_{11}^2}{N_1 p_{11} + N_2 p_{21}} + \frac{\beta_2 p_{12}^2}{N_1 p_{12} + N_2 p_{22}} \right) \frac{N_1}{\alpha_1} \left( \frac{\beta_1 p_{11} p_{21}}{N_1 p_{11} + N_2 p_{21}} + \frac{\beta_2 p_{12} p_{22}}{N_1 p_{12} + N_2 p_{22}} \right) \frac{N_2}{\alpha_2} \\ \left( \frac{\beta_1 p_{11} p_{21}}{N_1 p_{11} + N_2 p_{21}} + \frac{\beta_2 p_{12} p_{22}}{N_1 p_{12} + N_2 p_{22}} \right) \frac{N_1}{\alpha_1} \left( \frac{\beta_1 p_{21}^2}{N_1 p_{11} + p_{21} N_2} + \frac{\beta_2 p_{22}^2}{N_1 p_{12} + N_2 p_{22}} \right) \frac{N_2}{\alpha_2} \end{bmatrix}.$$

The relationship (13) is obtained by using the fact that, in (12), we have  $\dot{S}_i + \dot{I}_i = -\alpha_i I_i \leq 0$ . This implies that  $\lim_{t \rightarrow \infty} I_i(t) = 0$  (for  $i = 1, 2$ ), since  $S_i$  and  $I_i$  are positive and integrating  $\frac{\dot{S}_i}{S_i}$  in (12), we obtain, after some tedious algebra Expression (13). Brauer (2008) and Brauer and Castillo-Chávez (2012) give more details on the computation of the final size relationship. We would like to point out that the next-generation matrix and the matrix  $K$  defining the final epidemic size have the same eigenvalues.

The residence-time matrix  $\mathbb{P}$  plays an important role as evidenced by the dependence of the final epidemic size relation as in Fig. 7. As we can notice in Fig. 7, the prevalence in low-risk Patch 1 is highest in the high-mobility scheme, whereas in high-risk Patch 2, the high mobility leads to the lowest prevalence. Also, as stated before ( $\lim_{t \rightarrow +\infty} I_i(t) = 0$ , for  $i = 1, 2$ ) with any typical outbreak model, the disease ultimately dies out from both patches (Hethcote 2000).

### 5 Conclusion and Discussion

Heterogeneous mixing in multi-group epidemic models is most often defined in terms of group-specific susceptibility and average contact rates captured multiplicatively by the transmission parameter  $\beta$ . However, contact rates, in general, cannot be measured in satisfactory ways for diseases like influenza, measles or tuberculosis, due to the difficulty of assessing the average number of contacts per unit of time of susceptible populations in different locations for varied activities. In this paper, we propose the use of residence times in heterogeneous environments, as a proxy for “effective” contacts over a certain time window, and develop a multi-group epidemic framework via virtual dispersal where the risk of infection is a function of the residence time and local environmental risk. This novel approach eliminates the need to define and measure contact rates that are used in the traditional multi-group epidemic models with heterogeneous mixing.

Under the proposed framework, we formulate a general multi-patch SIS epidemic model with residence times. We calculate the basic reproduction number  $\mathcal{R}_0$  which is a function of a patch residence-time matrix  $\mathbb{P}$ . Our global analysis shows that the model is robust in the sense that the disease dynamics depend exclusively on the basic reproductive number when the residence-time matrix  $\mathbb{P}$  is “constant” (Theorem 2.1). We proved that the disease-free equilibrium is globally asymptotically stable (GAS) if the basic reproduction number  $\mathcal{R}_0 \leq 1$  and that a unique interior endemic GAS equilibrium exists if  $\mathcal{R}_0 > 1$ . This results holds as long as the residence-time matrix  $\mathbb{P}$  is irreducible; that is, the graph of the patches is strongly connected.

If the residence-time matrix  $\mathbb{P}$  is not irreducible, that is, the network of the patches is not strongly connected, Theorem 2.1 does not apply. For these cases, our further analysis (Theorem 2.2) provides accessible insights on the impact of the residence matrix  $\mathbb{P}$  on the infection levels within each patch. Our results imply that the infection risk (measured by  $\mathcal{B}$ ) and the residence-time matrix ( $\mathbb{P}$ ) can play an important role in the endemicity at the patch level. More specifically, the right combinations of the environmental risk level ( $\mathcal{B}$ ) and dispersal behavior ( $\mathbb{P}$ ) can either promote or suppress infection for particular patches. For example, we are able to apply Theorem 2.2 to the two-patch case when residents of Patch 1 visit Patch 2 but not conversely. Theorem 2.2 allows us not only to characterize the patch-specific disease dynamics as a function of the time spend by residents and visitors to the patch of interest, but also to classify patches as sources or sinks of infection, a role that depends on risk ( $\mathcal{B}$ ) and mobility ( $\mathbb{P}$ ).

The entries of residence-time matrix  $\mathbb{P}$  could be prevalence dependent, i.e., not constant anymore. The disease dynamics are expected to be different than the cases when  $\mathbb{P}$  is constant. To explore these differences, we study a two-patch model with the state-dependent residence-time matrix  $\mathbb{P}$ , and assume that each entry  $p_{ij}(I_1, I_2)$  of the residence times  $\mathbb{P}(I_1, I_2)$  is negatively correlated with the prevalence in Patch  $j$ . When the residence times  $\mathbb{P}(I_1, I_2)$  is prevalence dependent, our analysis and simulations suggest that (1) its disease dynamics may be prone to persistent by comparing to the case when  $\mathbb{P}$  is constant (e.g., Figs. 1, 2); and (2) the disease endemic level could be rather complicated (e.g., Figs. 3, 4, 5, 6).

We have extended our framework to a two-patch SIR single-outbreak model to explore how the residence-time matrix  $\mathbb{P}$  may affect the final endemic size. We first derived the final epidemic size relationship in order to capture the size of the outbreak. Our analysis and simulations support that the residence-time matrix  $\mathbb{P}$  plays an important role in the final epidemic size. For example, as observed in Fig. 7, the prevalence in low-risk Patch 1 is highest in the high-mobility scheme, whereas in high-risk Patch 2, the high mobility leads to the lowest prevalence.

In both conventional and phenomenological approaches to residence times used in this paper, humans behavior and responses to disease risk are automatic:  $\mathbb{P}$  is constant and predefined functions of health status. Recent studies (Fenichel et al. 2011; Horan and Fenichel 2007; Horan et al. 2011, 2010; Perrings et al. 2014) have incorporated behavior as a feedback response coupled with the dynamics of the disease. A model of the decision to spend time in patch  $i = 1, 2$  based on individuals' utility functions that include the possibility of adapting to changing contagion dynamics in the above two-patch setting, using previous work (Fenichel et al. 2011; Morin and Castillo-Chavez 2003), is the subject of a separate study.

**Acknowledgments** These studies were made possible by grant #1R01GM100471-01 from the National Institute of General Medical Sciences (NIGMS) at the National Institutes of Health. The contents of this manuscript are solely the responsibility of the authors and do not necessarily represent the official views of DHS or NIGMS. Research of Y.K. is partially supported by NSF-DMS (1313312). The funders had no role in study design, data collection and analysis, decision to publish or preparation of the manuscript. The authors are grateful to two anonymous referees for helpful comments and suggestions which led to an improvement of this paper.

### Appendix 1: Computation of $\mathcal{R}_0$

*Proof* The general SIS model with residence time is described by the system (6)

$$\dot{I} = \text{diag}(\tilde{N} - I)\mathbb{P}\text{diag}(\mathcal{B})\text{diag}(\tilde{N})^{-1}\mathbb{P}^t I - \text{diag}(d_I + \gamma_I)I.$$

The right-hand member of the above system can be clearly decomposed as  $\mathcal{F} + \mathcal{V}$  where

$$\mathcal{F} = \text{diag}(\tilde{N} - I)\mathbb{P}\text{diag}(\mathcal{B})\text{diag}(\tilde{N})^{-1}\mathbb{P}^t I \quad \text{and} \quad \mathcal{V} = -\text{diag}(d_I + \gamma_I)I$$

The Jacobian matrix at the DFE of  $\mathcal{F}$  and  $\mathcal{V}$  is given by:

$$F = D\mathcal{F}\Big|_{DFE} = \text{diag}(\tilde{N})\mathbb{P}\text{diag}(\mathcal{B})\text{diag}(\tilde{N})^{-1}\mathbb{P}^t \quad \text{and} \quad V = D\mathcal{V}\Big|_{DFE} = -\text{diag}(d_I + \gamma_I)$$

The basic reproduction number  $\mathcal{R}_0$  is given by the spectral radius of the next-generation matrix  $-FV^{-1}$  (Diekmann et al. 1990; Driessche and Watmough 2002). Hence, we deduce that

$$\mathcal{R}_0 = \rho(-\text{diag}(\tilde{N})\mathbb{P}\text{diag}(\mathcal{B})\text{diag}(\tilde{N})^{-1}\mathbb{P}^t V^{-1})$$

□

### Appendix 2: Proof of Theorem 2.1

The proof uses the method in Iggidr et al. (2012) which is based on Hirsch’s theorem (Hirsch 1984).

**Theorem 5.1** (Hirsch 1984) *Let  $\dot{x} = F(x)$  be a cooperative differential equation for which  $\mathbb{R}_+^n$  is invariant, the origin is an equilibrium, each  $DF(x)$  is irreducible, and that all orbits are bounded. Suppose that*

$$x > y \implies DF(x) < DF(y) \quad \text{for all } x, y.$$

*Then, all orbits in  $\mathbb{R}_+^n$  tend to zero or there is a unique equilibrium  $p^*$  in the interior of  $\mathbb{R}_+^n$  and all orbits in  $\mathbb{R}_+^n$  tend to  $p^*$ .*

*Proof of Theorem 2.1* Equation (6) can be written as:

$$\dot{I} = (F + V)I - \text{diag}(I)\mathbb{P}\text{diag}(\mathcal{B})\text{diag}(\tilde{N})^{-1}\mathbb{P}^t I \tag{14}$$

where  $F = \text{diag}(\tilde{N})\mathbb{P}\text{diag}(\mathcal{B})\text{diag}(\tilde{N})^{-1}\mathbb{P}^t$  and  $V = -\text{diag}(d_I + \gamma_I)$ , as defined in “Appendix 1.” Let us denote by  $X(I)$  the semi-flow induced by (14). Hence,

$$DX(I) = \text{diag}(\tilde{N} - I)\mathbb{P}\text{diag}(\mathcal{B})\text{diag}(\tilde{N})^{-1}\mathbb{P}^t + V - W(I_1, I_2) \tag{15}$$

where  $W(I_1, I_2) = \text{diag}(\mathbb{P}\text{diag}(\mathcal{B})\text{diag}(\tilde{N})^{-1}\mathbb{P}'I)$ . Since  $\mathbb{P}$  is irreducible and  $I \leq \tilde{N}$ ,  $DX(I)$  is clearly Metzler irreducible matrix. That means, the flow is strongly monotone. Plus,  $DX(I)$  is clearly decreasing with respect of  $I$ . Hence, by Hirsch's theorem all trajectories either go to zero or go to an equilibrium point  $\bar{I} \gg 0$ . From the relation (15), we have  $DX(0) = F + V$  where  $F$  and  $V$  are the one defined previously in "Appendix 1." However, since  $F$  a nonnegative matrix and  $V$  is Metzler, we have the following equivalence

$$\alpha(F + V) < 0 \iff \rho(-FV^{-1}) < 1$$

where  $\alpha(F + V)$  is the stability modulus, i.e., the largest real part of eigenvalues, of  $F + V$  and  $\rho(-FV^{-1})$  the spectral radius of  $-FV^{-1}$ . Hence, the DFE is globally asymptotically stable if  $\mathcal{R}_0 = \rho(-FV^{-1}) < 1$ . And if  $\mathcal{R}_0 > 1$ , i.e.,  $\alpha(F + V) > 0$ , the DFE is unstable (Driessche and Watmough 2002). Since, we have proved that  $DX(I)$  is a Metzler matrix, to prove the local stability of the endemic equilibrium  $\bar{I} \gg 0$ , we only need to prove that it exists  $w \gg 0$  such that  $DX(\bar{I})w < 0$  (Berman and Plemmons 1994). The endemic equilibrium  $\bar{I} \gg 0$  satisfies the equation

$$(F + V)\bar{I} - \text{diag}(\bar{I})\mathbb{P}\text{diag}(\mathcal{B})\text{diag}(\tilde{N})^{-1}\mathbb{P}'\bar{I} = 0$$

Hence,

$$DX(\bar{I})\bar{I} = -W(\bar{I})\bar{I} < 0$$

Hence, with  $w = \bar{I}$ , we deduce that  $\bar{I}$  is locally stable. With the attractivity of  $\bar{I}$  guaranteed Hirsh's theorem, we conclude that the endemic equilibrium  $\bar{I} \gg 0$  is globally asymptotically stable if  $\mathcal{R}_0 > 1$ .

Finally, if  $\mathcal{R}_0 = 1$ , we have  $\alpha(F + V) = 0$ . It exists  $c \gg 0$  such that  $(F + V)^t c = 0$ . By considering the Lyapunov function  $V = \langle c|I \rangle$ . This function is definite positive and its derivation along the trajectories if (14) is

$$\begin{aligned} \dot{V} &= \langle c|\dot{I} \rangle \\ &= \left\langle c|(F + V)I - \text{diag}(I)\mathbb{P}\text{diag}(\mathcal{B})\text{diag}(\tilde{N})^{-1}\mathbb{P}'I \right\rangle \\ &= -\left\langle c|\text{diag}(I)\mathbb{P}\text{diag}(\mathcal{B})\text{diag}(\tilde{N})^{-1}\mathbb{P}'I \right\rangle \\ &\leq 0 \end{aligned} \tag{16}$$

Plus  $\dot{V} = 0$  only at the DFE. Hence, the DFE is GAS if  $\mathcal{R}_0 = 1$ . This completes the proof of the Theorem 2.1.  $\square$

### Appendix 3: Proof of Theorem 2.2

*Proof* Since System (6) has an attracting compact  $\Omega$ , then according to Theorem (2.1), we can expect that  $\lim_{t \rightarrow \infty} I_i(t) < \frac{b_i}{d_i}$ ; thus, for time large enough, we can have  $\frac{b_i}{d_i} - I_i > 0$ , therefore we have

$$\dot{I}_i > I_i \left( \frac{b_i}{d_i} - I_i \right) \left( \sum_{j=1}^n \frac{\beta_j p_{ij}^2}{\sum_{k=1}^n p_{kj} \frac{b_k}{d_k}} \right) - (d_i + \gamma_i) I_i$$

which indicates that when  $\mathcal{R}_0^i(\mathbb{P}) > 1$

$$\dot{I}_i|_{I_i=0} = \frac{b_i}{d_i} \left( \sum_{j=1}^n \frac{\beta_j p_{ij}^2}{\sum_{k=1}^n p_{kj} \frac{b_k}{d_k}} \right) - (d_i + \gamma_i) > 0.$$

Then apply the average Lyapunov Theorem (Hutson 1984), we can conclude that  $\liminf_{t \rightarrow \infty} I_i(t) > 0$ ; i.e., the disease in the residence Patch  $i$  is persistent if  $\mathcal{R}_0^i(\mathbb{P}) > 1$ .

If  $p_{ij} > 0$  and  $p_{kj} = 0$  for all  $k = 1, \dots, n$ , and  $k \neq i$ , this implies that if there is a portion of the residence Patch  $i$  population flowing into the residence Patch  $j$ , then there is no other residence Patch  $k$  where  $k \neq j$ , i.e.,

$$\beta_j p_{ij} \sum_{k=1, k \neq i}^n p_{kj} I_k = 0$$

which also implies that

$$\left( \frac{b_i}{d_i} - I_i \right) \sum_{j=1}^n \frac{\beta_j p_{ij} \sum_{k=1, k \neq i}^n p_{kj} I_k}{\sum_{k=1}^n p_{kj} \frac{b_k}{d_k}} = 0.$$

then we can conclude that Model (6) can have an equilibrium since under these conditions,

$$\frac{b_i}{d_i} \sum_{j=1}^n \frac{\beta_j p_{ij} \sum_{k=1, k \neq i}^n p_{kj} I_k}{\sum_{k=1}^n p_{kj} \frac{b_k}{d_k}} = \frac{b_i}{d_i} \frac{\beta_i \sum_{k=1, k \neq i}^n p_{ki} I_k}{\sum_{k=1}^n p_{kj} \frac{b_k}{d_k}} = 0.$$

Therefore, if the conditions  $p_{kj} = 0$  for all  $k = 1, \dots, n$ , and  $k \neq j$  whenever  $p_{ij} > 0$  hold, then we have

$$\begin{aligned} \dot{I}_i|_{I_i=0} &= \left[ I_i \left( \frac{b_i}{d_i} - I_i \right) \left( \sum_{j=1}^n \frac{\beta_j p_{ij}^2}{\sum_{k=1}^n p_{kj} \frac{b_k}{d_k}} \right) \right. \\ &\quad \left. + \left( \frac{b_i}{d_i} - I_i \right) \sum_{j=1}^n \frac{\beta_j p_{ij} \sum_{k=1, k \neq i}^n p_{kj} I_k}{\sum_{k=1}^n p_{kj} \frac{b_k}{d_k}} - (d_i + \gamma_i) I_i \right] \Big|_{I_i=0} \\ &= \frac{b_i}{d_i} \sum_{j=1}^n \frac{\beta_j p_{ij} \sum_{k=1, k \neq i}^n p_{kj} I_k}{\sum_{k=1}^n p_{kj} \frac{b_k}{d_k}} = 0. \end{aligned}$$

Therefore,  $I_i = 0$  is the invariant manifold for Model (6).

On the other hand, when these conditions hold, then we have

$$\mathcal{R}_0^i(\mathbb{P}) = R_0^i \times \sum_{j=1}^n \left( \frac{\beta_j}{\beta_i} \right) p_{ij} \left( \frac{\left( p_{ij} \frac{b_i}{d_i} \right)}{\sum_{k=1}^n p_{kj} \frac{b_k}{d_k}} \right) = R_0^i \times \sum_{j=1}^n \left( \frac{\beta_j}{\beta_i} \right) p_{ij}.$$

Therefore, if  $\mathcal{R}_0^i(\mathbb{P}) = R_0^i \times \sum_{j=1}^n \left( \frac{\beta_j}{\beta_i} \right) p_{ij} < 1$ , then we have the following inequality:

$$\begin{aligned} \frac{\dot{I}_i}{I_i} &= I_i \left( \frac{b_i}{d_i} - I_i \right) \left( \sum_{j=1}^n \frac{\beta_j p_{ij}^2}{\sum_{k=1}^n p_{ki} \frac{b_k}{d_k}} \right) - (d_i + \gamma_i) I_i \\ &\leq I_i \left[ \frac{b_i}{d_i} \left( \sum_{j=1}^n \frac{\beta_j p_{ij}^2}{\sum_{k=1}^n p_{ki} \frac{b_k}{d_k}} \right) - (d_i + \gamma_i) \right] \\ &= I_i \left[ \sum_{j=1}^n \beta_j p_{ij} - (d_i + \gamma_i) \right] < 0. \end{aligned}$$

Therefore, we have  $\lim_{t \rightarrow \infty} I_i(t) = 0$ ; i.e., there is no endemic in the residence Patch  $i$ .  $\square$

## References

- Anderson RM, May RM (1982) Directly transmitted infectious diseases: control by vaccination. *Science* 215:1053–1060
- Anderson RM, May RM (1991) Infectious diseases of humans. Dynamics and control. Oxford Science Publications, New York
- Arino J (2009) Diseases in metapopulations. In: Ma Z, Zhou Y, Wu J (eds) Modeling and dynamics of infectious diseases, vol 11. World Scientific, Singapore
- Arino J, Davis J, Hartley D, Jordan R, Miller J, van den Driessche P (2005) A multi-species epidemic model with spatial dynamics. *Math Med Biol* 22:129–142
- Arino J, van den Driessche P (2003) The basic reproduction number in a multi-city compartmental model. *Lecture notes in control and information science*, vol 294, pp 135–142
- Arino J, van den Driessche P (2006) Disease spread in metapopulations. In: Zhao X-O, Zou X (eds) Nonlinear dynamics and evolution equations, vol 48. Fields Institute Communications, AMS, Providence, pp 1–13
- Berman A, Plemmons RJ (1994) Nonnegative matrices in the mathematical sciences, vol 9 of classics in applied mathematics. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA. Revised reprint of the 1979 original
- Bernoulli D (1766) Essai d'une nouvelle analyse de la mortalité causée par la petite vérole. *Mem. Math. Phys. Acad. R. Sci. Paris*, pp. 1–45
- Blythe SP, Castillo-Chavez C (1989) Like-with-like preference and sexual mixing models. *Math Biosci* 96:221–238
- Brauer F (2008) Epidemic models with heterogeneous mixing and treatment. *Bull Math Biol* 70:1869–1885
- Brauer F, Castillo-Chavez C (1994) Basic models in epidemiology. In: Steele J, Powell T (eds) Ecological time series. Raven Press, New York, pp 410–477

- Brauer F, Castillo-Chávez C (2012) Mathematical models in population biology and epidemiology. In: Marsden JE, Sirovich L, Golubitski M (eds) Applied mathematics, vol 40. Springer, New York
- Brauer F, Castillo-Chavez C, Velasco-Hernández JX (1996) Recruitment effects in heterosexually transmitted disease models. In: Kirschner D (ed) Advances in mathematical modeling of biological processes, vol 3:1. Int J Appl Sci Comput, pp 78–90
- Brauer F, Feng Z, Castillo-Chavez C (2010) Discrete epidemic models. *Math Biosci Eng* 7:1–15
- Brauer F, van den Driessche P (2001) Models for transmission of disease with immigration of infectives. *Math Biosci* 171:143–154
- Brauer F, van den Driessche P, Wang L (2008) Oscillations in a patchy environment disease model. *Math Biosci* 215:1–10
- Brauer F, Watmough J (2009) Age of infection epidemic models with heterogeneous mixing. *J Biol Dyn* 3:324–330
- Castillo-Chavez C, Busenberg S (1991) A general solution of the problem of mixing of subpopulations and its application to risk-and age-structured epidemic models for the spread of AIDS. *Math Med Biol* 8:1–29
- Castillo-Chavez C, Cooke K, Huang W, Levin SA (1989) Results on the dynamics for models for the sexual transmission of the human immunodeficiency virus. *Appl Math Lett* 2:327–331
- Castillo-Chavez C, Hethcote H, Andreasen V, Levin S, Liu W (1989) Epidemiological models with age structure, proportionate mixing, and cross-immunity. *J Math Biol* 27:233–258
- Castillo-Chavez C, Huang W (1999) Age-structured core group model and its impact on STD dynamics. In: *Mathematical approaches for emerging and reemerging infectious diseases: models, methods, and theory* (Minneapolis, MN, 1999), vol. 126 of IMA, Math Appl, Springer, New York, 2002, pp. 261–273
- Castillo-Chavez C, Huang W, Li J (1996) Competitive exclusion in gonorrhea models and other sexually transmitted diseases. *SIAM J Appl Math* 56:494–508
- Castillo-Chavez C, Huang W, Li J (1999) Competitive exclusion and coexistence of multiple strains in an SIS STD model. *SIAM J Appl Math* 59:1790–1811 (electronic)
- Castillo-Chavez C, Thieme HR (1995) Asymptotically autonomous epidemic models. In: Arino ADE, Kimmel O, Kimmel M (eds) *Mathematical population dynamics: analysis of heterogeneity, volume one: theory of epidemics*. Wuerz, Winnipeg
- Castillo-Chavez C, Velasco-Hernández JX, Fridman S (1994) Modeling contact structures in biology. In: Levin SA (ed) *Frontiers in mathematical biology*, vol 100. Springer, Berlin ch. 454–491
- Chowell D, Castillo-Chavez C, Krishna S, Qiu X, Anderson KS (2015) Modelling the effect of early detection of Ebola. *Lancet* 15:148–149
- Diekmann O, Heesterbeek JAP, Metz JAJ (1990) On the definition and the computation of the basic reproduction ratio  $R_0$  in models for infectious diseases in heterogeneous populations. *J Math Biol* 28:365–382
- Dietz K, Heesterbeek J (2002) Daniel Bernoulli's epidemiological model revisited. *Math Biosci* 180:1–21
- Dietz K, Schenzle D (1985) Mathematical models for infectious disease statistics. In: Atkinson, Anthony, Fienberg, Stephen E (eds) *A celebration of statistics*. Springer, New York, pp 167–204
- Fall A, Iggidr A, Sallet G, Tewa J-J (2007) Epidemiological models and Lyapunov functions. *Math Model Nat Phenom* 2:62–68
- Fenichel E, Castillo-Chavez C, Ceddia MG, Chowell G, Gonzalez Parra P, Hickling GJ, Holloway G, Horan R, Morin B, Perrings C, Springborn M, Valazquez L, Villalobos C (2011) Adaptive human behavior in epidemiological models. *PNAS* 108(15):6306–6311
- Guo H, Li M, Shuai Z (2006) Global stability of the endemic equilibrium of multigroup models. *Can Appl Math Q* 14:259–284
- Guo H, Li M, Shuai Z (2008) A graph-theoretic approach to the method of global Lyapunov functions. *Proc Am Math Soc* 136(8):2793–2802
- Hadeler K, Castillo-Chavez C (1995) A core group model for disease transmission. *Math Biosci* 128:41–55
- Heiderich KR, Huang W, Castillo-Chavez C (2002) Nonlocal response in a simple epidemiological model. In: *Appl IVM (ed) Mathematical approaches for emerging and reemerging infectious diseases: an introduction*, vol 125. Springer, New York, pp 129–151
- Hernandez-Ceron N, Feng Z, Castillo-Chavez C (2013) Discrete epidemic models with arbitrary stage distributions and applications to disease control. *Bull Math Biol* 75:1716–1746
- Hethcote HW (1976) Qualitative analyses of communicable disease models. *Math Biosci* 28:335–356
- Hethcote HW (2000) The mathematics of infectious diseases. *SIAM Rev* 42:599–653 (electronic)
- Hethcote HW, Thieme HR (1985) Stability of the endemic equilibrium in epidemic models with subpopulations. *Math Biosci* 75:205–227

- Hethcote HW, Yorke J (1984) *Gonorrhoea: transmission dynamics and control*, vol 56. Lecture notes in biomathematics. Springer
- Hirsch M (1984) The dynamical system approach to differential equations. *Bull AMS* 11:1–64
- Horan DR, Fenichel EP (2007) Economics and ecology of managing emerging infectious animal diseases. *Am J Agric Econ* 89:1232–1238
- Horan DR, Fenichel EP, Melstrom RT (2011) Wildlife disease bioeconomics. *Int Rev Environ Resour Econ* 5:23–61
- Horan DR, Fenichel EP, Wolf CA, Graming BM (2010) Managing infectious animal disease systems. *Annu Rev Resour Econ* 2:101–124
- Hsu Schmitz S-F (2000a) Effect of treatment or/and vaccination on HIV transmission in homosexual with genetic heterogeneity. *Math Biosci* 167:1–18
- Hsu Schmitz S-F (2000b) A mathematical model of HIV transmission in homosexuals with genetic heterogeneity. *J Theor Med* 2:285–296
- Hsu Schmitz S-F (2007) The influence of treatment and vaccination induced changes in the risky contact rate on HIV transmission. *Math Popul Stud* 14:57–76
- Huang W, Cooke K, Castillo-Chavez C (1992) Stability and bifurcation for a multiple-group model for the dynamics of HIV/AIDS transmission. *SIAM J Appl Math* 52:835–854
- Huang W, Cooke KL, Castillo-Chavez C (1992) Stability and bifurcation for a multiple-group model for the dynamics of HIV/AIDS transmission. *SIAM J Appl Math* 52:835–854
- Hutson V (1984) A theorem on average Lyapunov functions. *Monatshefte für Mathematik* 98:267–275
- Iggidr A, Sallet G, Tsanou B (2012) Global stability analysis of a metapopulation SIS epidemic model. *Math Popul Stud* 19:115–129
- Jacquez JA, Simon CP (1993) The stochastic SI model with recruitment and deaths I. Comparison with the closed SIS model. *Math Biosci* 117:77–125
- Jacquez JA, Simon CP, Koopman J (1991) The reproduction number in deterministic models of contagious diseases. *Comment Theor Biol* 2:159–209
- Jacquez JA, Simon CP, Koopman J, Sattenspiel L, Perry T (1988) Modeling and analyzing HIV transmission: the effect of contact patterns. *Math Biosci* 92:119–199
- Kuniya T, Muroya Y (2014) Global stability of a multi-group SIS epidemic model for population migration. *DCDS Ser B* 19(4):1105–1118
- Lajmanovich A, Yorke J (1976) A deterministic model for gonorrhoea in a nonhomogeneous population. *Math Biosci* 28:221–236
- Lin X, So JW-H (1993) Global stability of the endemic equilibrium and uniform persistence in epidemic models with subpopulations. *J Aust Math Soc Ser B* 34:282–295
- Morin B, Castillo-Chavez C (2003) SIR dynamics with economically driven contact rates. *Nat Resour Model* 26:505–525
- Mossong J, Hens N, Jit M, Beutels P, Mikolajczyk R, Massari M, Salmaso S, Tomba GS, Wallinga J, Heijne J, Sadkowska-Todys M, Rosinska M, Edmunds WJ (2008) Social contacts and mixing patterns relevant to the spread of infectious diseases. *PLoS Med* 5:381–391
- Nold A (1980) Heterogeneity in disease-transmission modeling. *Math Biosci* 52:227
- Perrings C, Castillo-Chavez C, Chowell G, Daszak P, Fenichel EP, Finnoff D, Horan RD, Kilpatrick AM, Kinzig AP, Kuminoff NV, Levin S, Morin B, Smith KF, Springborn M (2014) Merging economics and epidemiology to improve the prediction and management of infectious disease. *Ecohealth* 11(4):464–475
- Ross R (1911) *The prevention of malaria*. John Murray, London
- Rushton S, Mautner A (1955) The deterministic model of a simple epidemic for more than one community. *Biometrika* 42(1/2):126–132
- Sattenspiel L, Dietz K (1995) A structured epidemic model incorporating geographic mobility among regions. *Math Biosci* 128:71–91
- Sattenspiel L, Simon CP (1988) The spread and persistence of infectious diseases in structured populations. *Math Biosci* 90:341–366 [Nonlinearity in biology and medicine (Los Alamos, NM, 1987)]
- Shuai Z, van den Driessche P (2013) Global stability of infectious disease models using Lyapunov functions. *SIAM J Appl Math* 73:1513–1532
- Simon CP, Jacquez JA (1992) Reproduction numbers and the stability of equilibria of SI models for heterogeneous populations. *SIAM J Appl Math* 52:541–576
- van den Driessche P, Watmough J (2002) Reproduction numbers and sub-threshold endemic equilibria for compartmental models of disease transmission. *Math Biosci* 180:29–48

- Velasco-Hernández JX, Brauer F, Castillo-Chavez C (1996) Effects of treatment and prevalence-dependent recruitment on the dynamics of a fatal disease. *IMA J Math Appl Med Biol* 13:175–192
- Vidyasagar M (1980) Decomposition techniques for large-scale systems with nonadditive interactions: stability and stabilizability. *IEEE Trans Autom Control* 25:773–779
- Yorke JA, Hethcote HW, Nold A (1978) Dynamics and control of the transmission of gonorrhea. *Sex Transm Dis* 5:51–56